Etiopathogenesis of nasal polyps

Chojnowska S.^{1*}, Kępka A.², Waszkiewicz N.³, Kołodziejczyk ZP.⁴, Konarzewska-Duchnowska E.⁵, Ościłowicz K.⁴, Cabaj-Wiater I.⁶, Sowizdraniuk J.⁷, Dzięcioł J.⁸, Ładny JR.⁵, Zwierz K.⁴, Szajda SD.⁵

³ Department of Psychiatry, Medical University of Bialystok, Poland

⁴ Medical College of the Universal Education Society, Łomża, Poland

⁶ District Hospital of Garwolin, Poland

⁷ Emergency Service of Krakow, Poland

ABSTRACT

Nasal polyps are soft, pink or pearl white consensus among physic outgrowths usually connected with nasal mucosa by etiopathogenesis of nasa

flaccid crus. A histological image of nasal polyps indicates significant differences in particular cases. Because of the variety of nasal polyps, the choice of proper laryngological treatment is very difficult. Difficulties in treatment are due to the lack of

consensus among physicians regarding the etiopathogenesis of nasal polyps. Probably etiopathogenesis of nasal polyps is not homogeneous and polyps' formation is influenced by many coexisting mechanisms.

Key words: Nasal polyps, histology, treatment

¹ Medical Institute, College of Computer Science & Business Administration in Łomża, Poland

² Department of Biochemistry and Experimental Medicine, The Children's Memorial Health Institute in Warsaw, Poland

⁵ Department of Emergency Medicine and Disasters, Medical University of Bialystok, Poland

⁸ Department of Human Anatomy, Medical University of Bialystok, Poland